

Research Article

Cite this article: Schiønning MK, Marcús L, Veríssimo A, Thomasberger A, Ballyram SA, Försterra G and Concha FJ (2025) Sharks and skates in the Comau Fjord, northern Patagonia, Chile: an elasmobranch species checklist with biological aspects. *Journal of the Marine Biological Association of the United Kingdom* **105**, e44, 1–16. <https://doi.org/10.1017/S0025315425000335>

Received: 19 July 2024

Revised: 27 February 2025

Accepted: 2 March 2025

Keywords:

biodiversity, elasmobranchs, fjords, molecular analysis, nursery, Patagonia, phylogenetics, shark, skate, species inventory

Corresponding author: Mette K. Schiønning;
Email: mekjs@aqu.dtu.dk

Sharks and skates in the Comau Fjord, northern Patagonia, Chile: an elasmobranch species checklist with biological aspects

Mette K. Schiønning^{1,2} , Lara Marcús³, Ana Veríssimo^{4,5}, Aris Thomasberger^{1,2}, Stacy A. Ballyram^{2,6}, Günter Försterra^{2,6} and Francisco J. Concha^{7,8} 

¹Technical University of Denmark, National Institute of Aquatic Resources, Kgs. Lyngby, Denmark; ²Fundación San Ignacio del Huinay, Hualaihue, Puerto Montt, Chile; ³Facultad de Medicina y Ciencia, Universidad San Sebastian, Puerto Montt, Chile; ⁴CIBIO-InBIO, Research Center in Biodiversity and Genetic Resources, University of Porto, Vairão, Portugal; ⁵BIOPOLIS Program in Genomics, Biodiversity and Land Planning, CIBIO, Vairão, Portugal; ⁶Escuela de Ciencias del Mar, Facultad de Recursos Naturales, Pontificia Universidad Católica de Valparaíso, Valparaíso, Chile; ⁷Facultad de Ciencias del Mar y de Recursos Naturales, Universidad de Valparaíso, Valparaíso, Chile and ⁸Department of Ecology and Evolutionary Biology, University of Connecticut, Storrs, CT, USA

Abstract

Fjord ecosystems serve as crucial habitats for elasmobranchs, supporting them across all life stages. Chilean Patagonia provides one of the most complex and extensive networks of fjord ecosystems in the world, displaying high marine biodiversity, including elasmobranchs. However, little is known about this ecologically important group of fishes in these ecosystems. This study investigates the biodiversity of elasmobranchs in the Comau Fjord over a period of 6 months by combining morphological and molecular data. In total, 309 specimens within a radius of 7.5 km were recorded, belonging to six families and nine species: *Hexanchus griseus* (77.5–178 cm L_T), *Notorynchus cepedianus* (180.6 cm L_T), *Schroederichthys bivius* (35–65.2 cm L_T), *Scymnodon macracanthus* (37.3 cm L_T), *Centrophorus squamosus* (87.4 cm L_T), *Deania calceus* (58.3–98.6 cm L_T), *Squalus acanthias* (25.5–101.1 cm L_T), *Dipturus chilensis* (62.9–152 cm L_T), and *Dipturus trachyderma* (69.8–194 cm L_T). This included records of three species previously unknown in the fjord and was equivalent to nearly 20% of the elasmobranch richness found in Southern Chile. The results further suggest that the Comau Fjord could be a primary nursery ground for several species of elasmobranchs. This is the first time that a species inventory of elasmobranchs is conducted in a Chilean fjord system. The outcomes of this research provide an elasmobranch species checklist with biological aspects from the Comau Fjord, which are essential data to inform decision makers, conservation managers, and future research.

Introduction

Fjord ecosystems are important habitats for elasmobranchs (sharks, rays, and skates) across all of their life stages, functioning as feeding, nursing, and pupping grounds (Andrews et al., 2010; Gaitán-Espitia et al., 2017; Licandeo and Cerna, 2007). The Chilean Patagonian fjords comprise a large geographic area of ~240,000 km² and more than 84,000 km of coastline, with a complex network of unique ecosystems and high marine species richness (Friedlander et al., 2021; Mutschke, 2008). Nonetheless, many of the fjords in northern Chilean Patagonia, especially those in the Chiloense Ecoregion (see Spalding et al., 2007) have been impacted by anthropogenic stressors for decades, including the development of the salmon aquaculture industry and overfishing (Buschmann et al., 2023; Molinet and Niklitschek, 2023). Growing economy and population, as well as stressors related to climate change, are likely to further increase the pressure on the fjords' ecosystem and its biodiversity (e.g. Soto et al., 2019, 2023).

Elasmobranchs are a diverse group of cartilaginous fishes with important ecological functions (Heithaus et al., 2022, 2010) and generally slow, complex life history strategies (Ebert et al., 2021; Last et al., 2016), which make them highly susceptible to anthropogenic and environmental disturbances (Dulvy et al., 2021; Ferretti et al., 2010; Pacoureau et al., 2021). The most recent checklist of elasmobranchs in Chile included 101 species, of which nearly half of them occur within the latitudes of Southern Chile (c. 36° S to c. 54° S), including fjord ecosystems (Bustamante et al., 2014). Largely due to little direct economic importance, most elasmobranchs have only received minor research attention in the Chilean fjords, resulting in a lack of information on their conservation status and functional role within these ecosystems (Molinet and Niklitschek, 2023).

© The Author(s), 2025. Published by Cambridge University Press on behalf of Marine Biological Association of the United Kingdom. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.



In contrast to most of the fjords in Chilean Patagonia, the Comau Fjord has been subject to extensive scientific research during the past 20 years (e.g. Försterra and Häussermann, 2003; Villalobos et al., 2021). While the research has primarily focused on the sessile benthic fauna of the area, few studies have been dedicated to the ichthyofauna. Unpublished data from 2007 (N. Straube, personal communication, May 2, 2023) and data from 2013 located on the online Global Cestode Database (Caira et al., 2024) show that at least six species of elasmobranchs are present in the Comau Fjord, namely spiny dogfish *Squalus acanthias* (Squalidae), narrowmouthed catshark *Schroederichthys biviis* (Atelomycteridae), bluntnose sixgill shark *Hexanchus griseus* (Hexanchidae), leafscale gulper shark *Centrophorus squamosus* (Centrophoridae), birdbeak dogfish *Deania calceus* (Centrophoridae), and roughskin skate *Dipturus trachyderma* (Rajidae).

Considering the existing knowledge gap due to a lack of elasmobranch-related research, and the potential threats to the fjord ecosystems in northern Patagonia, elasmobranch species inventories and information about their life history stages, abundance, seasonality, and migration are necessary to provide decision makers with the knowledge needed to counteract potential local extirpations and to conserve biodiversity. The present study combines morphological and molecular data: (i) to provide a species checklist with images of confirmed reports in the Comau Fjord and (ii) to gain insights into the biological aspect of the examined specimens (i.e. sex, length, and maturity) and their relative abundances. These data are further discussed in relation to the fjord acting as an elasmobranch habitat.

Materials and methods

Study area

The field survey took place in the Comau Fjord (42°15' S, 72°29' W), which is located in northern Chilean Patagonia (Figure 1). The region is characterised by deep and narrow fjords carved by ice masses during the last glacial period. The Comau Fjord is 2–5 km wide and stretches over 40 km from its head in the south to its mouth in the north. The depth at the centre of the fjord ranges from approximately 300 m at its head to around 500 m close to its mouth (Addamo et al., 2021). Freshwater and glacial meltwater influx from terrestrial runoff creates a pronounced low salinity surface layer, which varies in its bathymetric extension between < 1 m in summer and up to 10 m in winter, with strong local, seasonal, and weather-dependant changes in extension, temperature (5–20 °C), and salinity (2–20 PSU), followed by a brackish water layer (20–30 PSU, 12–20 °C) at 1–20 m (Sánchez et al., 2011). These low-salinity surface and brackish subsurface layers are especially pronounced close to the head of the fjord where the highest freshwater input occurs (Addamo et al., 2021). The conditions become fully marine below 20 m, with a year-round stable salinity of >32 PSU and temperatures of 10.2–12.0 °C (Fillinger and Richter, 2013; Sánchez et al., 2011). Tidal currents with amplitudes of up to 7.5 m in combination with wind driven currents can, however, on occasions mix water masses down to 100 m depth (Addamo et al., 2021). As documented from other fjord regions around the world, the Comau Fjord also exhibits features of deepwater emergence, which is a phenomenon referring to the occurrence of deepwater species in shallow depths, far beyond their usual distributional range (Feehan et al., 2019; Fillinger and Richter, 2013). While the drivers beyond this phenomenon are not yet fully

understood (Häussermann et al., 2021), its occurrence could be an indication for bathyal- or abyssal-like conditions in the study area, thereby providing refugia to deepwater and cold-adapted organisms.

Sampling of specimens

Specimens ($n = 305$) were sampled from May to October 2018 at 10 locations (Figure 1) with demersal longlines, soaked overnight for 14–16 h at a range of depths 80–450 m. The groundline was held down with weights and fitted with circle hooks set 10 m apart. Buoys were placed between the weights to keep the groundline off the seabed to minimise depredation by epibenthic crustaceans. The hooks were baited with different species of fish, such as Chilean silverside (*Odontesthes regia*), Patagonian grenadier (*Macruronus magellanicus*), and other miscellaneous whitefish. In addition, four elasmobranchs that were opportunistically sampled and/or recorded from January 2017 to April 2018 were included in the study (Figure 1, Locations 11–14). These samples comprised two whole specimens donated to the authors (Figure 1, Locations 11 & 14), one *in situ* personal observation (Figure 1, Location 12) and another one as photographic evidence (Figure 1, Location 13).

All specimens were identified in the field to species level following the diagnostic characters proposed by Ebert (2015) and Ebert and Mostarda (2016). They were further photographed with a NIKON 3300 camera to build up a photo voucher collection and to support taxonomic identification. When possible, both total length (L_T) and clasper length (L_C in males) of the individuals were measured down to the nearest 1 mm. Sex was determined based on the presence or absence of claspers. The maturity of males was determined by the degree of calcification of the claspers and their length in relation to the pelvic fins based on the reproductive stage protocols by ICES (2010). The degrees of maturity are: (I) immature males: non-calcified claspers shorter than the pelvic fins; (II) developing males: partially calcified claspers extending to the margin of the pelvic fins or slightly beyond, and (III) mature males: fully calcified and long claspers extending well beyond the margin of the pelvic fins. In this study, reproductive stages of females could not be determined since most of the specimens were alive and subsequently released after capture to keep the mortality rate at the minimum. On a few dead female specimens ($n = 4$), the ovaries were, however, examined for oocytes using reproductive stage protocols and relevant literature (Hamlett WC, 2005; ICES, 2010). Individuals with visible vitelline scars (i.e. fresh or healed) were also noted (see Castro, 1993).

Molecular analysis

Tissue samples were collected from specimens ($n = 273$) and were immediately preserved in 96% ethanol and frozen at –20 °C until further molecular analysis. Genomic DNA (gDNA) was extracted using commercial kits EasySpin® Genomic DNA Tissue Kit (Citomed) following the manufacturer's recommendations and checked for quality on 0.8% agarose gel electrophoresis run at 300 V with 0.5X TBE buffer. Nucleotide sequences of the mitochondrial ND2 gene were obtained via the polymerase chain reaction (PCR) using the primers ASNM and ILEM by Naylor et al. (2012). All samples were amplified in 10 µl reactions, including 5 µl of MyTaq HS Master mix, 3 µl of autoclaved water, 0.4 µl of each primer (10 µM) and 1 µl of gDNA. The PCR temperature profile included an initial denaturation step of 5 min at 95 °C, followed by 30 cycles of denaturation for 1 min at 94 °C, annealing for 30 sec




Figure 1. Map of the Comau Fjord (42°S) with the sampling locations (A) within South America (B) and northern Patagonia in Chile (C). The monthly sampling of the elasmobranchs was conducted from May to October 2018 on locations 1–10. The additional locations 11–14 represent the opportunistically sampled and observed specimens from January 2017 to April 2018. The contour lines are expressed as depth in meters (50–450 m depth).

at 55 °C, and extension for 1 min at 72 °C, and by a final extension step of 10 min at 60 °C. Successful PCR amplification was checked on 2% agarose gel electrophoresis and the obtained amplicons were enzymatically cleaned with ExoSAP-IT (Thermo Fisher Scientific) prior to Sanger sequencing on a commercial service provider. No successful amplification of the ND2 gene was obtained for the specimens morphologically identified as *Hexanchus griseus*, thus, we performed molecular identification using the MiFish-E primers of Miya et al. (2015), which amplify a small fragment (~172 bp) of the mitochondrial 12S ribosomal RNA gene (12S). The same PCR protocol was used as described above, but the temperature profile included an initial denaturation step of 3 min at 95 °C, followed by a touchdown of nine steps of 95 °C for 15 sec, 54 °C for 10 sec (decreasing 0.5 °C per cycle), and 72 °C for 10 sec, followed by 31 cycles of 95 °C for 5 sec, 50 °C for 10 sec, and 72 °C for 10 sec, plus a final extension step at 72 °C for 5 min. The amplicons were

cleaned with ExoSAP-IT and processed for Sanger sequencing, as described above.


The obtained nucleotide sequences were checked for quality and manually edited in Geneious Prime (version 2020.1.2). Additional ND2 sequences from a subset of species (i.e. those that were successfully amplified as described above) were obtained from Naylor et al. (2012), Straube et al. (2013), and Concha et al. (2019) referring to specimens collected in the corresponding type locality (Table 1). The type localities and taxonomic status of all species were checked following Fricke et al. (2016).

Comparison of the newly generated sequences to those of the putative species' type locality was performed by aligning all sequences using the ClustalW algorithm (Thompson et al., 1994) as implemented in Geneious Prime, and by constructing the respective neighbour-joining tree using *p*-distances and 1000 bootstrap replicates in MEGA version X (Stecher et al., 2020). For

Table 1. Reference sequences of the ND2 genes used in molecular identification and phylogenetic analyses

Species	Type locality	GenBank Accession no.	Voucher locality
<i>Hexanchus griseus</i>	Mediterranean, France	JQ518727	Eastern Atlantic
<i>Notorynchus cepedianus</i>	Tasmania, Australia	JQ518728	South Australia (Great Bight)
<i>Schroederichthys bivi</i>	Cape of Good Hope*	JQ518684	Argentina
<i>Centrophorus squamosus</i>	Eastern North Atlantic	JQ518947	Madeira, Portugal
<i>Deania calceus</i>	Madeira, Portugal	JQ519007	Southwestern Indian
<i>Squalus acanthias</i>	Mediterranean	JQ518978	Rhode Island, USA
<i>Dipturus chilensis</i>	Valparaíso, Chile	MK613955	Valparaíso, Chile
<i>Dipturus trachyderma</i>	Southern Argentina	OR813835	Argentina

*Locality stated in original type description is likely an error as the species is endemic to South America (Compagno, 1984; Springer, 1979).

**Figure 2.** Stacked bar chart shows the relative species composition at each sampling station (1–10) and the opportunistically sampled and observed specimens (11–14).

the specific case of putative *H. griseus* specimens, the obtained sequences were compared against those deposited in GenBank, using the BLASTn algorithm (Camacho *et al.*, 2009).

Results

Relative species composition and abundance

In this study, 309 elasmobranch specimens were morphologically identified in the Comau Fjord, comprising nine species of six families. These comprised three new species records in the fjord, namely broadnose sevengill shark (*Notorynchus cepedianus*), largespine velvet dogfish (*Scymnodon macracanthus*), and yellownose skate (*Dipturus chilensis*). The most abundant species were spiny dogfish (*Squalus acanthias*), narrowmouthed catshark (*Schroederichthys bivi*), and yellownose skate (*Dipturus chilensis*), with the numbers caught, length ranges (by sex), and the proportional occurrence in sampling sites, as summarised in Table 2 and Figure 2.

Molecular analysis

Nucleotide sequences of the ND2 gene (1034 bp) were generated for 15 specimens (GenBank Accession no.s PV268444 to

PV268458), comprising all the species identified based on morphology, except *S. macracanthus*, for which no tissue was available. The resulting neighbour-joining tree, including newly generated sequences from a few representative individuals per species and reference sequences from type locality specimens of each species (where available), shows congruence in morphological and molecular species identification (Figure 3). Nucleotide sequences of the 12S gene (172 bp) were generated for seven specimens, morphologically identified as *H. griseus* (GenBank Accession no. PV275033), which were 100% similar. BLASTn results showed 99% identity (83% query coverage) with other *H. griseus* sequences available on GenBank of unknown origin and with a specimen from Suruga Bay, Japan (NC_022732.1), supporting congruence between morphological and molecular identification.

Family Hexanchidae

Hexanchus griseus (Bonnaterre, 1788)

Bluntnose sixgill shark (English), cañabota gris (Spanish) (Figures 3 and 4A–E)

Type locality: France, northwestern Mediterranean Sea.

A total of 14 *H. griseus* specimens were recorded, comprising five females and nine immature males (Table 2 and Figure 2). Twelve specimens were processed for molecular identification with

Table 2. Results overview showing by family; species; sex; total number (*n*) of specimens; relative species composition of the longlined specimens (%); range of total lengths (cm), life stages as immature (I), developing (II) and mature (III), in the Comau Fjord from 2017 to 2018. The estimated length-at-50%-maturity (L_m) for females of each species is shown under the table

Family	Species	Sex	Abundance		Range (cm)	Maturity		
			<i>n</i>	%		I	II	III
Hexanchidae	<i>Hexanchus griseus</i>	Total	14	4.6	77.5–178			
		Male	9		77.5–178	9	0	0
		Female ^a	5		78.8–115			
		Unknown	0					
	<i>Notorynchus cepedianus</i>	Total	3		180.6			
		Male	0					
		Female ^b	1		180.6		1	
		Unknown	2					
Atelomycteridae	<i>Schroederichthys bivius</i>	Total	30	9.8	35–65.2			
		Male	28		35–65.2	1	11	16
		Female ^c	2		45.4–47.9			
		Unknown	0					
Somniosidae	<i>Scymnodon macracanthus</i>	Total	1	0.3				
		Male	1		37.3			
		Female ^d	0					
		Unknown	0					
Centrophoridae	<i>Centrophorus squamosus</i>	Total	1	0.3				
		Male	1		87.4			1
		Female ^f	0					
		Unknown	0					
	<i>Deania calceus</i>	Total	14	4.3	58.3–98.6			
		Male	4		71–75.3	0	1	3
		Female ^e	10		59.1–98.6			3
		Unknown	0					
Squalidae	<i>Squalus acanthias</i>	Total	214	70.2	25.5–101.1			
		Male	156		29.7–76.2	17	5	134
		Female ^g	41		25.5–101.1			
		Unknown	17					
Rajidae	<i>Dipturus chilensis</i>	Total	28	9.2	62.9–152			
		Male	4		67.1–86.2	0	2	2
		Female ^h	23		62.2–152			
		Unknown	1					
	<i>Dipturus trachyderma</i>	Total	4	1.3				
		Male	1		194			1
		Female ⁱ	2		69.8–157			
		Unknown	1					

^a L_m 421 cm (Ebert, 1986).

^b L_m 190 cm (Irigoyen et al., 2018).

^c L_m 48.5 cm (Colonello et al., 2020).

^d L_m 170 cm (Ebert et al., 2021).

^e L_m 85 cm (Lamilla and Bustamante, 2005).

^f L_m 119 cm (Parker and Francis, 2012).

^g L_m 72.8 cm (Gaitán-Espitia et al., 2017).

^h L_m 103 cm (Licandeo and Cerna, 2007).

ⁱ L_m 215 cm (Licandeo et al., 2007).




Figure 3. Neighbour-joining tree of *p*-distances among nucleotide sequences of the ND2 gene. Bootstrap support values were generated using 1000 replicates and are shown on the branches if >75%. Reference sequences are shown with GenBank Accession numbers, while newly generated sequences are shown with the sample number after species name. No sequence from *Dipturus trachyderma* was included in the analysis since the sequence length obtained was much shorter than those from other taxa.




Figure 4. *Hexanchus griseus* (Hexanchidae) specimens. (A) Lateral view of an immature female (#46). (B) Ventral view of the head of an immature male (#229). (C) Portion of the upper set of teeth (#114). (D) Portion of the lower set of teeth. (E) Young juvenile with a visible, healed vitelline scar (#228).

the ND2 genes but none amplified the target fragment. Indeed, we confirmed that the ILEM primer had no complementarity to the *H. griseus* ND2 gene. We have therefore used a small fragment of the 12S gene (~172 bp) for the same purpose. Seven sequences were successfully obtained, all of which matched *H. griseus* sequences with >99% identity against the NCBI *nt* database (NCBI Resource Coordinators, 2024).

The colouration of the dorsal side of the examined specimens was grey to dark grey or dark brown and occasionally blackish. They only comprised one dorsal fin located near the caudal area. The specimens had a light-coloured lateral line extending from about the middle of the trunk into the caudal fin. The colouration of the ventral surface differed among the specimens, being slightly lighter ventrally on the smaller individuals to darker and more uniform on the larger specimens. The head was broad and blunt, with a wide mouth, as reported by Compagno (1984). The most useful characteristics for identifying *H. griseus* from the regional relatives was by counting the pairs of gill openings, since it comprises six pairs of gill openings as opposed to seven pairs in *N. cepedianus* and *Heptranchias perlo*. Moreover, the *H. griseus* specimens did not have small black dots, as noticed on the *N. cepedianus* specimen (see next section).

Notorynchus cepedianus (Péron, 1807)

Broadnose sevengill shark (English), tiburón vaca de hocico corto (Spanish)

(Figures 3 and 5A–G)

Type locality: Adventure Bay, Tasmania, Australia.

Two of the three *Notorynchus cepedianus* specimens were observed in March and December 2017 (Table 2 and Figure 2).

The single sampled specimen (September 2017) was a developing female that contained small, visible oocytes. The observed specimens were likely of a similar size. The single sample included in the analysis showed 99% similarity with a reference sequence from South Australia (Great Bight) and formed a well-supported clade (Figure 3).

The dorsal coloration of the examined specimen was grey to dark grey with many black spots and some sporadic white spots, as reported by Compagno (1984) and Ebert et al. (2021). The dorsal side comprised one dorsal fin near the caudal area. The colouration of the ventral side was light to white. The head was broad and rounded with a wide mouth, as reported by Compagno (1984) and Ebert et al. (2021). There are a number of useful diagnostic features of *N. cepedianus* that can be used to distinguish it from the related species *H. griseus* and *H. perlo*. However, since two of the specimens were never physically sampled and only exist as photographic evidence, counting the pairs of gill openings and examining the morphology of the teeth and the head was not possible. Instead, the observed specimens were distinguished from the two regional relatives based on the presence of the sporadic white spots on the dorsal side, which are absent in both *H. griseus* and *H. perlo*, and are a distinct feature of *N. cepedianus* only, in the family Hexanchidae (Compagno, 1984; Ebert et al., 2021). Apart from the presence of the spots, the sampled specimen was distinguished from *H. griseus* by the additional pair of gill openings. As for *H. perlo*, which is a much smaller species reaching a maximum size of about 140 cm L_T (Ebert et al., 2021), the exceedingly narrow head and the morphological features of the teeth were used as characteristics to distinguish it from the *N. cepedianus* specimen. As of 2024, the number of observations of *N. cepedianus* is only increasing and is spotted in



Figure 5. *Notorynchus cepedianus* (Hexanchidae) specimens. (A) Lateral view of a developing female specimen (#6). (B) Dorsal view of the head and the trunk. (C) Ventral view of the head. (D) A portion of the upper set of teeth. (E) A portion of the lower set of teeth. (F) Screenshot of a video with an *N. cepedianus* encounter during a SCUBA dive at 10 meters depth in December 2017 (Figure 1, Location 12). (G) An *N. cepedianus* caught near a salmon farm in March 2017 by a local fisher (Figure 1, Location 13).

much shallower depths (B. Hernández, personal communication, 23 March 2024).

Family Atelomycteridae

Schroederichthys bivius (Smith, 1838)

Narrowmouthed catshark (English), pintarroja (Spanish) (Figures 3 and 6A–D)

Type locality: Cape of Good Hope (but likely an error; see Springer (1979) and Compagno (1984)).

In total, 30 *S. bivius* specimens were recorded during the study, which comprised two females and 28 males (Table 2 and Figure 2). Four samples were sequenced for the ND2 gene and formed a single, well-supported clade with the reference sequences from the southwestern Atlantic (Figure 3).

The dorsal surfaces of *S. bivius* specimens were brownish with multiple brown and white dots and large rhombic reddish to dark brown patches along the body. The patches had small white and dark reddish spots. The ventral surface of the specimens was whitish and lacked spots. *S. bivius* specimens were distinguished from the regional congener *S. chilensis*, based on distinct diagnostic features (e.g. saddles), following the taxonomic description by Compagno (1984).

Family Somniosidae

Scymnodon macracanthus (Regan, 1906)

Largespine velvet dogfish (English), tollo, sapata espinuda (Spanish) (Figures 3 and 7A–C)

Type locality: Strait of Magellan, Chile.

One immature male *S. macracanthus* was recorded but not sampled during the study (Table 2 and Figure 2). Thus, this species was only morphologically identified, based on several diagnostic features that distinguishes it from other Somniosidae species in the Southeastern Pacific Ocean.

The colouration of the specimen was dark brown to black, and it had relatively large spiracles. The snout was moderately long and rounded. The dorsal fins, which had a prominent and somewhat large dorsal spine, were greater in length than in height. The second dorsal fin was taller than the first (Ebert *et al.*, 2021).

Family Centrophoridae

Centrophorus squamosus (Bonnaterre, 1788)

Leafscale gulper shark (English), quelvacho negro (Spanish) (Figures 3 and 8A–C)

Type locality: not stated (probably eastern North Atlantic).

A single *Centrophorus squamosus* specimen was recorded in this study, which was a developing male (Table 2 and Figure 2). Genetically, the examined specimen showed 100% similarity to the reference sequence from the Eastern North Atlantic, off Madeira, Portugal (Figure 3).

The colour of the specimen was plain dark brownish to black. Its snout was moderately flattened, with a preoral snout length shorter than the distance between the mouth and pectoral fins origin. The dorsal fins were about the same height, but the first dorsal fin was more elongated. Both fins had prominent dorsal spines. This species is similar to birdbeak shark (*Deania calceus*). However, the preoral snout length (i.e. shorter in *C. squamosus*) and interdorsal space (i.e. longer in *C. squamosus*) were useful characteristics to ensure a more precise identification.

Deania calceus (Lowe, 1839)

Birdbeak dogfish (English), tollo pajarito (Spanish) (Figures 3 and 9A–D)

Type locality: Madeira, Portugal, eastern Atlantic.

A total of 14 *D. calceus* specimens were recorded, comprising 10 females and four males (Table 2 and Figure 2). The four samples sequenced for the ND2 gene formed a single, highly supported clade with a reference sequence from the Southwestern Indian Ocean (Figure 3).

The colouration of the dorsal side of the examined *D. calceus* specimens was dark grey and brown to black and slightly lighter ventrally. They were distinguished from morphologically similar species in the region, such as *C. squamosus*, by its relatively (as proportion of L_T) longer preoral snout length and relatively (as proportion of L_T) shorter interdorsal space (see Ebert, 2015).

Family Squalidae

Squalus acanthias Linnaeus, 1758

Spiny dogfish/spurdog (English), tollo de cacho (Spanish) (Figures 3 and 10A–D)

Type locality: Mediterranean Sea and northeastern Atlantic.



Figure 6. *Schroederichthys bivius* (Atelomycteridae) specimens. (A) Lateral view of a developing male (#156). (B) Dorsal view of the head and the trunk of a developing male (#8). (C) Ventral view of the head and the trunk of a mature male (#174). (D) Lateral view of the head and the trunk of a mature male (#256).



Figure 7. *Scymnodon macracanthus* (Somniosidae) specimen (#239). (A) Dorsal view. (B) Dorsal view of the head. (C) Ventral view of the head and the trunk.




Figure 8. *Centrophorus squamosus* (Centrophoridae) specimen (#69). (A) Lateral view of a mature male. (B) Ventral view of the head. (C) Dorsal view of the head and the trunk.




Figure 9. *Deania calceus* (Centrophoridae) specimens. (A) Lateral view of a female (#217). (B) Ventral view of the head of a female (#218). (C) Dorsal view of the head (#218). (D) Upper and lower set of the teeth of a mature female (#121).

In total, 214 *S. acanthias* specimens were recorded, comprising 41 females, 156 males, and 17 unsexed and unmeasured specimens (Table 2 and Figure 2). About a quarter (24%) of all the measured specimens comprised juveniles (<50 cm) and among these, at least seven specimens had visible, translucent vitelline scars (22.5–40.5 cm L_T). A single sample was sequenced for the

ND2 gene and showed 99% similarity to a reference sequence from Rhode Island, USA, forming a well-supported clade (Figure 3).

The dorsal sides of examined specimens of *S. acanthias* were light to dark grey with some or several conspicuous white spots on their flanks. The colouration laterally faded towards the ventral side. These specimens had a moderately long and pointed snout,




Figure 10. *Squalus acanthias* (Squalidae) specimens. (A) Lateral view of an immature male (#133). (B) Dorsal view of the head of a mature male (#146). (C) Ventral view of the head and the trunk of a mature male (#128). (D) Immature male with a healed translucent vitelline scar (#5).

and a markedly larger second dorsal fin spine, closely to extending beyond the dorsal fin on some specimens. The ventral surface of the specimens was white with the occasional darker pigmented patterns. The underside of the pectoral fins was darker than the remaining dorsal side.

Family Rajidae

Dipturus chilensis (Guichenot, 1848)

Yellownose skate (English), raya volántin (Spanish)
(Figures 3 and 11A–D)

Type locality: Valparaíso Bay, central Chile, southeastern Pacific Ocean.

A total of 28 *D. chilensis* specimens were recorded, comprising 23 females, four males, and one unsexed and unmeasured specimen (Table 2 and Figure 2). The four specimens of *D. chilensis* included in the molecular analysis showed >99% of similarity with the sequence data of the neotype from Valparaíso, Chile, (GenBank Accession no. MK613955; Concha *et al.*, 2019) and clustered in the same, highly supported clade (Figure 3).

The dorsal surface of the examined specimens was dark brownish, with randomly distributed lighter spots, and a large red to purple ocellum near the centre of each pectoral fin. The ventral surface of the body was dark, almost as dark as the dorsal surface and certainly darker than what was reported from central Chile (see Concha *et al.*, 2019). The colouration patterns on the dorsal surface (i.e. lighter spots and purple to reddish ocelli) were variable, compared to the morphologically similar *D. trachyderma* specimens. These were mostly plain grey to blackish and lacked coloured ornamentation. In addition, the spinulation pattern of the caudal fin on *D. chilensis* specimens contained up to three rows of caudal thorns, as opposed to *D. trachyderma* specimens, which could comprise up to six rows.

Dipturus trachyderma (Krefft & Stehmann, 1975)

Roughskin skate (English), Raya espinosa (Spanish).
(Figures 3 and 12A–D)

Type locality: Off southern Argentina, southwestern Atlantic




Figure 11. *Dipturus chilensis* (Rajidae) specimens. (A) Dorsal view of a female (#223). (B) Ventral view of a developing male (#150). (C) Dorsal view of a female with visible ocelli on the pectoral fins (#238). (D) Dorsal view of the head of a female with lighter spots (#223).




Figure 12. *Dipturus trachyderma* (Rajidae) specimens. (A) Dorsal view of a female (#22). (B) Ventral view of a female (#22). (C) Dorsal view of the head of a large mature male (#240). (D) Ventral view of pointy claspers (#240).

In this study, four specimens of *D. trachyderma* were collected, comprising two females, one mature male, and one unmeasured and unsexed specimen (Table 2 and Figure 2). One specimen of *D. trachyderma* included in the molecular analysis showed 100% identity to the reference sequence from off Argentina, near the type locality (OR813835; Table 1). However, given the smaller length of the reference sequence (590 bp) compared to the ND2 sequences obtained here, *D. trachyderma* was not included in the neighbour-joining tree.

Lacking any colour patterns, both dorsal and ventral side of *D. trachyderma* specimens were dark brownish, with a slightly lighter ventral side, as originally described by Krefft and Stehmann (1975) and reported by Leible and Stehmann (1987). The spinulation was restricted to the caudal area, which comprised several rows of caudal thorns, as opposed to *D. chilensis* specimens (see previous section). The size, colouration, and spinulation were the most useful characters in identifying this species from its sympatric congener, *D. chilensis*.

Discussion

This study presents a species checklist of elasmobranchs inhabiting the Comau Fjord in northern Patagonia (42° S) by combining morphological identification with molecular analysis to validate species identification. It provides valuable new information about the elasmobranch biodiversity in the Comau Fjord and reveals that it is equivalent to nearly 20% of the species richness found in Southern Chile (Bustamante et al., 2014), with at least 55% being threatened with extinction (Table 3). A number of studies from Southern Chile have focused on the life history traits and ecology of a few species but not the entire elasmobranch community, hence, community-focused studies on elasmobranchs are still scarce in the region (see Gaitán-Espitia et al., 2017; Licandeo and Cerna, 2007; Vargas-Caro et al., 2017). Therefore, this is to the best of our knowledge the first species inventory of elasmobranchs in a Chilean fjord system.

Based on a sample size of 309 specimens, the study recorded nine species of elasmobranchs (i.e. seven species of shark and two species of skate), which includes three species not previously

documented to inhabit the Comau Fjord (i.e. broadnose sevengill shark *Notorynchus cepedianus*, largespine velvet dogfish *Scymnodon macracanthus*, and yellownose skate *Dipturus chilensis*; Table 3). The most abundant species was spiny dogfish (*Squalus acanthias*; 70.2%) with the next most abundant species being narrowmouthed catshark (*Schroederichthys bivius*; 9.8%) and yellownose skate (*D. chilensis*; 9.2%), which are commonly known to be bycaught or targeted species in many small- to large-scale fisheries in southern Chile (De la Torre et al., 2010; Lamilla et al., 2010; Quiroz et al., 2008; Valenzuela et al., 2008). The examined specimens were consistent in both morphologic and molecular data when compared to other reference specimens or descriptions from other regions of the world. Consistency between morphological and genetic data of the specimens included in this study, when compared to the same species from type localities, became relevant in confirming taxonomic identity of the studied species.

Essential habitat: primary nursery ground

In this study, no neonates with fresh or unhealed scars were collected, yet, several young *S. acanthias* juveniles with visible, translucent vitelline scars were registered at the head of the fjord in late autumn (Figures 2 and 10D). Since neonates consume smaller food items such as shrimps and amphipods (Sulikowski et al., 2013), it is unlikely that they would bite the whole hooked baits deployed for sampling in this study, accounting for their absence in the sample population. *S. acanthias* neonates are about 18–33 cm L_T at birth (Ebert et al., 2021) and may grow 10–15 cm during the first years of their life (Holden and Meadows, 1964), which suggests that some of the young juveniles collected in the fjord should be young-of-the-year (YOY; Northeast Fisheries Science Center, 2006).

It is generally unknown whether parturition of *S. acanthias* occurs in enclosed, coastal or offshore waters in the eastern South Pacific, as detailed information about their biology and ecology in this geographic area is still lacking (De la Torre et al., 2010). Yet, other observations from the same biogeographic province found that nursery areas of *S. acanthias* appeared to be located

Table 3. A presence (+)/absence (–) list of elasmobranchs observed in the Comau Fjord in 2007, 2013, and 2017–2018, including their status on the IUCN Red List (i.e. EN = Endangered; VU = Vulnerable; NT = Near Threatened; LC = Least Concern; DD = Data Deficient) and a downwards arrow to indicate a status with a decreasing trend (↓)

Family	Species	Years			Status	Reference
		2007	2013	2017–2018		
Hexanchidae	<i>Hexanchus griseus</i>	-	+	+	NT↓	(Finucci et al., 2020b)
	<i>Notorynchus cepedianus</i>	-	-	+	VU↓	(Finucci et al., 2020a)
Atelomycteridae	<i>Schroederichthys bivius</i>	+	+	+	LC	(Dulvy et al., 2020b)
Somniosidae	<i>Scymnodon macracanthus</i>	-	-	+	DD	(Finucci and Kyne, 2018)
Centrophoridae	<i>Centrophorus squamosus</i>	+	-	+	EN↓	(Finucci et al., 2020c)
	<i>Deania calceus</i>	+	+	+	NT↓	(Finucci et al., 2020e)
Squalidae	<i>Squalus acanthias</i>	+	+	+	VU↓	(Finucci et al., 2020d)
Rajidae	<i>Dipturus chilensis</i>	-	-	+	EN↓	(Dulvy et al., 2020c)
	<i>Dipturus trachyderma</i>	+	-	+	EN↓	(Dulvy et al., 2020a)

in open waters of the Southwest Atlantic (Colonello et al., 2016). The migration range of immature *S. acanthias*, however, is thought to be rather limited (Hickling, 1930), while the onshore–offshore migrations should only occur with sexual maturity, typically as a response to the seasonal changes in the water temperature (Carlson et al., 2014; Compagno, 1984). A migration of neonates from an offshore area to the study sites in the inner part of the fjord shortly after birth would seem unlikely. Thus, the results obtained in this study could imply that an enclosed environment, such as the Comau Fjord, functions as primary nursery area (see Bass, 1978; Heupel et al., 2007) for *S. acanthias*.

The notion of the fjord serving as a primary nursery area could possibly extend to other shark species. All the *H. griseus* specimens were exclusively juveniles of mixed sex (≤ 180 cm L_T) and considerably far from reaching either adolescence or adulthood (Ebert, 1986; Table 2). *H. griseus* sharks are approximately 61–93 cm L_T at birth (Ebert, 1986, 2002) and may double their length within the first year of life (Ebert, 2003). In this study, some specimens were <90 cm L_T with translucent but healed vitelline scars and a lighter ventral side (Figure 4E), indicating that young juveniles frequent the fjord and that some could be YOY. It is not uncommon that neonate and juvenile sixgill sharks exhibit site fidelity and that they may occupy shallower inshore waters like estuaries, bays, and fjords (Andrews et al., 2007; Ebert, 2002; King and Cornthwaite (Née Surry), 2017; Williams et al., 2010). For example, in some fjord-like ecosystems such as the Strait of Georgia and the Puget Sound (northeast Pacific), *H. griseus* juveniles use the areas as nursery grounds before migrating offshore to the deeper continental slope as they mature (Andrews et al., 2007; King and Cornthwaite (Née Surry), 2017; Williams et al., 2010).

Segregation by sex and life history stages

Many elasmobranchs live in social units segregated by sex and size, divided into schools of mature males, mature females, and immature individuals of mixed sex (Klimley, 2013). In this study, there was a much lower frequency of mature *S. acanthias* females (1%) in relation to mature males (68%), and at three stations (i.e. 3, 6, and 10), there was a notable spatial overlap of immature individuals and mature males. This is in contrast to a previous study conducted in the nearby Chiloé Sea, which showed a high frequency of females

in all life history stages during summer and winter, with generally a low proportion of males (Gaitán-Espitia et al., 2017). Unlike this study, Gaitán-Espitia et al. (2017) also noticed a spatial overlap between mature females and immature individuals. It is well known that *S. acanthias* usually occurs in large schools with immature fish segregating by size and mature fish segregating by size and sex (Nammack et al., 1985; Stenberg, 2005; Templeman, 1944). This has even led to female-biased fish mortality in the north-west Atlantic (Haugen et al., 2017). Here, we can add that there also seems to be spatial segregation in southern Chilean waters, in addition to school segregation by size and sex. There are other studies suggesting similar spatial segregation in this species tied to environmental/abiotic factors (Dell’Apa et al., 2016; Sagarese et al., 2013), although male-avoidance behaviour in terms of avoiding courtship could also play an important role (Sims, 2006).

Segregation by size and sex has also been reported for *H. griseus* (Barnett et al., 2012; Ebert, 1989, 2002) and *S. bivius* (Colonello et al., 2020). The results from the current study imply an absence of mature *H. griseus* individuals in the Comau Fjord. While it must be acknowledged that such indications of size segregations could be due to sampling bias, it could also indicate a strategy to maximise survival, as previously hypothesised by King and Cornthwaite (Née Surry) (2017). Segregation by life history stages was similarly found in Southern Africa, where juvenile *H. griseus* sharks occupy shallower areas than adults, which likely minimises resource overlap, intraspecific competition, and the potential of cannibalism by adults (Barnett et al., 2012).

Although based on a limited sample size, there was a bias in the sex ratio of *S. bivius*, showing a higher frequency of males (93.3%) than females (6.7%), of which 64% of the males were mature. This could indicate spatial segregation by sex for this species in the Chilean Patagonia, as noted in the populations in Argentine Patagonia (see Colonello et al., 2020). As a sexually dimorphic species, with males maturing and attaining a larger size than the females (Cedrola et al., 2012), sexual segregation could occur due to the different physiological needs and energy requirements between the sexes (Colonello et al., 2020; Sims, 2006).

Limitations and future studies

The main limitation of this study is the scale of sampling. Due to resource constraints, sampling was only possible in two seasons of

the year. Furthermore, there were more opportunities for sampling in the southern area of the fjord, leading to a spatial sampling bias. As a result, seasonality in terms of spatial segregations among the species could not be evaluated. Moreover, some species may not have been represented due to eventual seasonality in their migration habits. In order to address such limitations, future systematic long-term spatial and temporal sampling covering the entire fjord, potentially in a combination with conventional tagging or acoustic telemetry could be initiated (e.g. Andrews et al., 2010; Carlson et al., 2014). This could elucidate how the elasmobranchs use the fjord as a habitat through time and space, any presence of other species, and how anthropogenic stressors might affect the distribution and abundance of this group of fishes. In addition, future studies should also consider including more non-invasive and cost-effective methods such as eDNA (environmental DNA) sampling and underwater camera traps to record the elasmobranch diversity and abundance in the fjord, as it is combined likely to have a higher species detection rate compared to other conventional sampling methods (Bakker et al., 2017; Boussarie et al., 2018; Liu et al., 2022). Furthermore, Local Ecological Knowledge (LEK) and sight reporting on social media could likely provide additional data on the presence/absence of the elasmobranchs and their estimated abundance and distribution, as another inexpensive and non-invasive method (see Batista et al., 2024).

While the current study has provided insights into the elasmobranch community of the Comau Fjord, effective conservation management of elasmobranchs relies on consistent and systematic monitoring of their biology and ecology. Elasmobranchs play important ecological roles in many ecosystems, both directly and indirectly. Thus, further research directed towards understanding their role in the fjord ecosystem and their resilience to anthropogenic stressors is warranted to ensure the long-term existence of the elasmobranch community in the Comau Fjord.

Data availability statement. The authors followed the style sheet found in <https://www.cambridge.org/core/journals/journal-of-the-marine-biological-association-of-the-united-kingdom/information/author-instructions/preparing-your-materials>

Acknowledgements. The authors are grateful for the help and assistance by Boris “Cormorán” and for sharing all his local knowledge and fishing expertise. A special thanks to Juan Pablo Espinoza, research assistant at Fundación San Ignacio del Huinay (FSIH), and all the volunteers at FSIH Stephanny, Curaz Leiva, Anna Covey, and Nicola Mohr, who assisted with the field sampling. Special thanks are extended to the Fundación San Ignacio del Huinay for their funding and support, which made this research possible.

Author contributions. Conceptualisation by M.K.S. and F.J.C.; fieldwork by M.K.S., A.T., and S.A.B.; methodology by M.K.S., A.T., and A.V.; analysis and interpretation of results by M.K.S., A.V., and F.J.C.; writing—original draft preparation by M.K.S., A.T., A.V., and F.J.C.; writing—review and editing by M.K.S., L.M., A.V., A.T., S.A.B., G.F., and F.J.C.; supervision by F.J.C. and L.M.; project administration by M.K.S.; sample logistics by L.M., F.J.C., and S.A.B.; funding acquisition by G.F. All authors have read and agreed to the published version of the manuscript.

Funding. M.K.S., A.T., and S.A.B. were financed by the Enel Generation Chile and the Pontificia Universidad Católica de Valparaíso, as the founding partners of Fundación San Ignacio del Huinay (FSIH), during the fieldwork. F.J.C. received support from Fulbright Chile, Faculty Development Program award, and from the Comisión Nacional de Investigación Científica y Tecnológica, CONICYT Becas Chile award. This work was supported in part by Dr. Janine N. Caira with funds from the National Science Foundation Awards, Nos. 1457762 and 1457776. A.V. was funded by Fundação para

a Ciência e Tecnologia (10.54499/DL57/2016/CP1440/CP1646/CT0001 and 10.54499/2022.02937.CEECIND/CP1730/CT0011).

Competing interests. The author(s) declare none.

Ethical standards. All elasmobranchs were carefully handled and sampled following the standard international and national protocol for handling marine life, in accordance with the sampling permit approved by the Subsecretaría de Pesca y Acuicultura, SUBPESCA (Resolución Exenta N° 1760).

References

- Addamo A, Zaccara S, Försterra G, Höfer J, García-Jiménez R, Crosa G and Machordom A (2021) Genetic conservation management of marine resources and ecosystems of Patagonian Fjords. *Frontiers in Marine Science* 8, 96.
- Andrews K, Levin P, Katz SL, Farrer D, Gallucci VF and Bargmann G (2007) Acoustic monitoring of sixgill shark movements in Puget sound: Evidence for localized movement. *Canadian Journal of Zoology* 85, 1136–1143.
- Andrews KS, Williams GD and Levin PS (2010) Seasonal and ontogenetic changes in movement patterns of sixgill sharks. *PLoS One* 5, e12549.
- Bakker J, Wangensteen OS, Chapman DD, Boussarie G, Buddo D, Guttridge TL, Hertler H, Mouillot D, Vigliola L and Mariani S (2017) Environmental DNA reveals tropical shark diversity in contrasting levels of anthropogenic impact. *Scientific Reports* 7, 16886.
- Barnett A, Braccini JM, Awruch CA and Ebert DA (2012) An overview on the role of Hexanchiformes in marine ecosystems: Biology, ecology and conservation status of a primitive order of modern sharks. *Journal of Fish Biology* 80, 966–990.
- Bass AJ (1978) Problems in studies of sharks in the Southwest Indian Ocean. Hodgson ES and Mathewson RF (eds), Arlington, Virginia, USA: Office of Naval Research, Department of the Navy, 545–594.
- Batista MI, Abril C, Verissimo A, Vasconcelos R, Pais M and Henriques S (2024) Mapping elasmobranch occurrences and overlap with human activities using local knowledge and non-invasive sampling to identify areas of potential conflict. *Frontiers in Marine Science* 11.
- Boussarie G, Bakker J, Wangensteen OS, Mariani S, Bonnin L, Juhel J-B, Kiszka JJ, Kulbicki M, Manel S, Robbins WD, Vigliola L and Mouillot D (2018) Environmental DNA illuminates the dark diversity of sharks. *Science Advances* 4, eaap9661. doi:10.1126/sciadv.aap9661
- Buschmann AH, Niklitschek EJ and Pereda SV (2023) Aquaculture and its impacts on the conservation of Chilean Patagonia. In Castilla JC, Zamudio JJA, Martínez-Harms MJ and Tecklin D (eds), *Conservation in Chilean Patagonia: Assessing the State of Knowledge, Opportunities, and Challenges*. Cham, Switzerland: Springer International Publishing, 303–320.
- Bustamante C, Vargas-Caro C and Bennett MB (2014) Not all fish are equal: Functional biodiversity of cartilaginous fishes (Elasmobranchii and Holocephali) in Chile. *Journal of Fish Biology* 85, 1617–1633.
- Caira JN, Jensen K and Barbeau E (2024) Global Cestode database. World Wide Web electronic publication. Available at www.elasmobranchs.tapewormdb.uconn.edu (accessed 17 June 2024).
- Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K and Madden TL (2009) BLAST+: Architecture and applications. *BMC Bioinformatics* 10, 421.
- Carlson AE, Hoffmayer ER, Tribuzio CA and Sulikowski JA (2014) The use of satellite tags to redefine movement patterns of spiny dogfish (*Squalus acanthias*) along the U.S. east coast: Implications for fisheries management. *PLoS One* 9, e103384.
- Castro J (1993) The shark nursery of Bulls Bay, South Carolina, with a review of the shark nurseries of the southeastern coast of the United States. *Environmental Biology of Fishes* 38, 37–48.
- Cedrola P, González A, Chiaramonte G and Pettovello A (2012) Bycatch of sharks (Elasmobranchii) in the Patagonian red shrimp *Pleoticus muelleri* (Bate, 1888) fishery. *Revista Del Museo Argentino de Ciencias Naturales, Nueva Serie* 14, 349–356.

- Colonello JH, Cortés F and Belleggia M (2020) Male-biased sexual size dimorphism in sharks: The narrowmouth catshark *Schroederichthys bivi* as case study. *Hydrobiologia* **847**, 1873–1886.
- Colonello JH, Cortés F, Belleggia M and Massa AM (2016) Reproductive and population parameters of spiny dogfish *Squalus acanthias* in the southwestern Atlantic Ocean. *Journal of Fish Biology* **88**, 1758–1775.
- Compagno LJV (1984) Sharks of the world. An annotated and illustrated catalogue of shark species known to date. Part 2 - Carcharhiniformes. *FAO Species Catalogue*, Vol. 4, 251–655. Rome, Italy: Food Aand Agriculture Organization Of The United Nations.
- Concha FJ, Caira JN, Ebert DA and Pompert JHW (2019) Redescription and taxonomic status of *Dipturus chilensis* (Guichenot, 1848), and description of *Dipturus lamillai* sp. nov. (Rajiformes: Rajidae), a new species of long-snout skate from the Falkland Islands. *Zootaxa* **4590**, 501–524.
- De la Torre A, Quinones R, Miranda D, Echevarría F, Quiñones and Urbina M (2010) South American sea lion and spiny dogfish predation on artisanal catches of southern hake in fjords of Chilean Patagonia. *ICES Journal of Marine Science* **67**, 294–303.
- Dell'Ápica A, Pennino M and Bonzek C (2016) Modeling the habitat distribution of spiny dogfish (*Squalus acanthias*), by sex, in coastal waters of the northeastern United States. *Fishery Bulletin* **115**, 89–100.
- Dulvy N, Acuña E, Bustamante C, Chiaramonte G, Concha F, Cuevas J, Herman K, Paesch L, Pompert J, Rincon G and Vélez-Zuazo X (2020a) *Dipturus trachydermus*, Roughskin Skate. In *The IUCN Red List of Threatened Species*. United Kingdom: International Union for Conservation of Nature and Natural Resources, 1–13.
- Dulvy N, Acuña E, Bustamante C, Chiaramonte G, Cuevas J, Herman K, Pompert J and Vélez-Zuazo X (2020b) *Schroederichthys bivi*, Narrowmouth Catshark. In *The IUCN Red List of Threatened Species*. United Kingdom: International Union for Conservation of Nature and Natural Resources, 1–9.
- Dulvy N, Acuña E, Bustamante C, Concha F, Herman K and Vélez-Zuazo X (2020c) *Dipturus chilensis*. In *The IUCN Red List of Threatened Species*. United Kingdom: International Union for Conservation of Nature and Natural Resources, 1–12.
- Dulvy NK, Pacoureau N, Rigby CL, Pollom RA, Jabado RW, Ebert DA, Finucci B, Pollock CM, Cheok J, Derrick DH, Herman KB, Sherman CS, VanderWright WJ, Lawson JM, Walls RHL, Carlson JK, Charvet P, Bineesh KK, Fernando D, Ralph GM, Matsushiba JH, Hilton-Taylor C, Fordham SV and Simpfendorfer CA (2021) Overfishing drives over one-third of all sharks and rays toward a global extinction crisis. *Current Biology* **31**, 4773–4787.
- Ebert D (1986) Biological aspects of the Sixgill Shark, *Hexanchus griseus*. *Copeia* **1986**, 131–135.
- Ebert D (1989) *The taxonomy, biogeography and biology of cow and frilled sharks (Chondrichthyes: Hexanchiformes)*. Department of Ichthyology and Fisheries Science. PhD Thesis, Rhodes University.
- Ebert D (2002) Some observations on the reproductive biology of The sixgill shark *Hexanchus griseus* (Bonnaterre, 1788) from Southern African waters. *South African Journal of Marine Science* **24**, 359–363.
- Ebert D (2003) *Sharks, Rays, Chimaeras of California*. University of California Press: California, USA.
- Ebert D (2015). In Mostarda E (Ed.), *Deep-sea Cartilaginous Fishes of the Southeastern Atlantic Ocean*. Rome, Italy: Food and Agriculture Organization.
- Ebert D, Dando M and Fowler S (2021) *Sharks of the World: A Complete Guide*. Princeton University Press: Princeton, USA.
- Ebert D and Mostarda E (2016) *Guía Para la Identificación de Peces Cartilaginosos de Aguas Profundas Del Océano Pacífico Sudoriental*. Food and Agriculture Organization - United Nations: Roma, Italia.
- Feehan KA, Waller RG and Häussermann V (2019) Highly seasonal reproduction in deep-water emergent *Desmophyllum dianthus* (Scleractinia: Caryophylliidae) from the Northern Patagonian Fjords. *Marine Biology* **166**, 1–13.
- Ferretti F, Worm B, Britten GL, Heithaus MR and Lotze HK (2010) Patterns and ecosystem consequences of shark declines in the ocean. *Ecology Letters* **13**, 1055–1071. doi:10.1111/j.1461-0248.2010.01489.x
- Fillinger L and Richter C (2013) Vertical and horizontal distribution of *Desmophyllum dianthus* in Comau Fjord, Chile: A cold-water coral thriving at low pH. *PeerJ* **1**.
- Finucci B, Barnett A, Cheok J, Cotton C, Kulka D, Neat F, Pacoureau N, Rigby C, Tanaka S and Walker T (2020a) *Notorynchus cepedianus*-Broadnose Sevengill Shark. In *The IUCN Red List of Threatened Species 2020*. United Kingdom: International Union for Conservation of Nature and Natural Resources, 1–11.
- Finucci B, Bineesh KK, Cheok J, Cotton C, Dharmadi D, Graham K, Kulka DW, Neat F, Pacoureau N, Rigby C, Tanaka S and Walker T (2020b) *Hexanchus griseus* - Bluntnose Sixgill Shark. In *The IUCN Red List of Threatened Species 2020*. United Kingdom: International Union for Conservation of Nature and Natural Resources, 1–13.
- Finucci B, Bineesh KK, Cheok J, Cotton CF, Dharmadi KDW, Neat FC, Pacoureau N, Rigby CL, Tanaka S and Walker TI (2020c) *Centrophorus squamosus*. In *The IUCN Red List of Threatened Species*. United Kingdom: International Union for Conservation of Nature and Natural Resources 1–16.
- Finucci B, Cheok J, Chiaramonte G, Cotton C, Dulvy N, Rigby C, Pacoureau N, Tanaka S and Walker T (2020d) *Squalus acanthias*. In *The IUCN Red List of Threatened Species*. United Kingdom: International Union for Conservation of Nature and Natural Resources, 1–18.
- Finucci B, Cheok J, Kulka D, Neat F, Pacoureau N, Rigby C, Tanaka S and Walker T (2020e) *Deania calcea* - Birdbeak Dogfish. In *The IUCN Red List of Threatened Species 2020*. United Kingdom: International Union for Conservation of Nature and Natural Resources, 1–13.
- Finucci B and Kyne PM (2018) *The IUCN Red List of Threatened Species Centroscyms macracanthus*, Largespine Velvet Shark. United Kingdom: International Union for Conservation of Nature and Natural Resources, 1–6.
- Försterra G and Häussermann V (2003) First report on large scleractinian (Cnidaria: Anthozoa) accumulations in cold-temperate shallow water of south Chilean fjords. *BMC Evolutionary Biology* **345**.
- Fricke R, Eschmeyer WN and der Laan R V (2016, April 1) Catalog of Fishes: Genera, Species, References. Available at <http://researcharchive.calacademy.org/research/ichthyology/catalog/fishcatmain.a> (accessed 16 June 2024).
- Friedlander AM, Ballesteros E, Goodell W, Hüne M, Muñoz A, Salinas-de-león P, Velasco-Charpentier C and Sala E (2021) Marine communities of the newly created Kawésqar National Reserve, Chile: From glaciers to the Pacific Ocean. *Plos One* **16**. doi:10.1371/journal.pone.0249413
- Gaitán-Espitia JD, Gómez D, Hobday AJ, Daley R, Lamilla J and Cárdenas L (2017) Spatial overlap of shark nursery areas and the salmon farming industry influences the trophic ecology of *Squalus acanthias* on the southern coast of Chile. *Ecology and Evolution* **7**, 3773–3783.
- Hamlett WC (2005) *Reproductive Biology and Phylogeny of Chondrichthyes: Sharks, Batoids, and Chimaeras*, Vol. 3. Boca Raton, Florida, USA: CRC Press.
- Haugen JB, Curtis TH, Fernandes PG, Sosebee KA and Rago PJ (2017) Sexual segregation of spiny dogfish (*Squalus acanthias*) off the northeastern United States: Implications for a male-directed fishery. *Fisheries Research* **193**, 121–128.
- Häussermann V, Ballyram S, Försterra G, Cornejo C, Ibáñez C, Sellanes J, Thomasberger A, Espinoza J and Beaujot F (2021) Species that fly at a higher game: Patterns of deep-water emergence along the Chilean Coast, including a global review of the phenomenon. *Frontiers in Marine Science* **8**.
- Heithaus M, Dunn R, Farabaugh N, Lester E, Madin E, Meekan M, Papastamatiou Y, Roff G, Vaudo J and Wirsing A (2022) *Advances in Our Understanding of the Ecological Importance of Sharks and Their Relatives*. Boca Raton, FL: CRC Press, 487–521.
- Heithaus M, Frid A, Vaudo J, Worm B and Wirsing A (2010) Unraveling the Ecological Importance of Elasmobranchs. In Jeffery C, John M and Michael H (eds), *Sharks and Their Relatives II: Biodiversity, Adaptive Physiology, and Conservation*. Boca Raton, Florida, USA: CRC Press, 611–637.
- Heupel MR, Carlson JK and Simpfendorfer CA (2007) Shark nursery areas: Concepts, definition, characterization, and assumptions. *Marine Ecology Progress Series* **337**, 287–297.
- Hickling CF (1930) A contribution towards the life-history of the Spur-dog. *Journal of the Marine Biological Association of the United Kingdom* **16**, 529–576. doi:10.1017/S0025315400072933

- Holden MJ and Meadows PS** (1964) The Fecundity of the Spurdog (*Squalus acanthias* L.). *ICES Journal of Marine Science* **28**, 418–424.
- ICES** (2010) Report of the Workshop on Sexual Maturity Staging of Elasmobranchs (WKMSSEL), p. 32. Valetta, Malta: International Council for the Exploration of the Sea, ICES Advisory Committee.
- Irigoyen A, De Wysiecki A, Trobbiani G, Bovcon N, Awruch C, Argemi F and Jaureguizar A** (2018) Habitat use, seasonality and demography of an apex predator in a marine temperate environment: The case of the sevengill shark *Notorynchus cepedianus* in northern Patagonia. *Marine Ecology Progress Series* **603**.
- King J and Cornthwaite (Née Surry) AM** (2017) Seasonal and daily movements of the bluntnose sixgill shark (*Hexanchus griseus*) in the strait of Georgia from satellite tag data. *Environmental Biology of Fishes* **100**.
- Klimley P** (2013) *The Biology of Sharks and Rays*. University of Chicago Press: Chicago, USA.
- Krefft G and Stehmann M** (1975) Results of the research trips of the FFS “Walther Herwig” to South America. XXXVI. Two further new ray species from the Southwest Atlantic: *Raja (Dipturus) leptocauda* and *Raja (Dipturus) trachyderma* spec. nov. (Chondrichthyes, Rajidae). *Archives of Fisheries Science* **26**, 77–97.
- Lamilla J and Bustamante C** (2005) *Guía Para El Reconocimiento De: Tiburones, Rayas Y Quimeras de Chile*. Santiago, Chile: Oceana.
- Lamilla J, Bustamante C, Roa R, Acuña E, Concha F, Meléndez R, López S, Aedo G, Flores H and Vargas-Caro C** (2010) *Estimación Del Descarte de Condrictios En Pesquerías Artesanales. Informe Final, Proyecto FIP N° 2008-60. Subsecretaría de Pesca*. Universidad Austral de Chile: Valdivia, Chile.
- Last P, Seret B, Stehmann M, Naylor G, White W, Carvalho M, White W, Carvalho M, Stehmann M and Naylor G** (2016) *Rays of the World*. CSIRO Publishing: Melbourne, Australia.
- Leible MD and Stehmann M** (1987) First records of *Raja (Dipturus) trachyderma* Krefft and Stehmann, 1975 from the southeastern Pacific off Chile, with first descriptions of its clasper characters and additional skeletal and morphological details (Pisces, Rajiformes, Rajidae). *Studies on Neotropical Fauna and Environment*, 169–188. doi:10.1080/01650528709360732
- Licandeo R, Cerna F and Céspedes R** (2007) Age, growth, and reproduction of the roughskin skate, *Dipturus trachyderma*, from the southeastern Pacific. *ICES Journal of Marine Science* **64**, 141–148. doi:10.1093/icesjms/fsl012
- Licandeo R and Cerna FT** (2007) Geographic variation in life-history traits of the endemic kite skate *Dipturus chilensis* (Batoidea: Rajidae), along its distribution in the fjords and channels of southern Chile. *Journal of Fish Biology* **71**, 421–440. doi:10.1111/j.1095-8649.2007.01499.x
- Liu Z, Collins R, Baillie C, Rainbird S, Brittain R, Griffiths A, Sims D, Mariani S and Genner M** (2022) Environmental DNA captures elasmobranch diversity in a temperate marine ecosystem. *Environmental DNA* **4**, 1024–1038. doi:10.1002/edn3.294
- Miya M, Sato Y, Fukunaga T, Sado T, Poulsen JY, Sato K, Minamoto T, Yamamoto S, Yamanaka H, Araki H, Kondoh M and Iwasaki W** (2015) MiFish, a set of universal PCR primers for metabarcoding environmental DNA from fishes: Detection of more than 230 subtropical marine species. *Royal Society Open Science* **2**. doi:10.1098/rsos.150088
- Molinet C and Niklitschek EJ** (2023). In Carlos CJ, J AZJ, José M-HM and David T (eds.), *Fisheries and Marine Conservation in Chilean Patagonia*. Cham, Switzerland: Springer International Publishing, 285–302.
- Mutschke E** (2008) Macrobenthic biodiversity and community structure in austral Chilean channels and fjords. In Nelson S and Sergio P (eds), *Progress in the Oceanographic Knowledge of Chilean Interior Waters, from Puerto Montt to Cape Horn*. Comité Oceanográfico Nacional-Pontificia Universidad Católica de Valparaíso. Valparaíso, Chile: National Oceanographic Committee, 133–141.
- Nammack M, Musick J and Colvocoresses J** (1985) Life history of Spiny Dogfish off the Northeastern United States. *Transactions of the American Fisheries Society* **114**, 367–376. doi:10.1577/1548-8659(1985)114<367:LHOSDO>2.0.CO;2
- Naylor G, Caira JN, Jensen K, Rosana KAM, White W and Last P** (2012) A DNA sequence-based approach to the identification of shark and ray species and its implications for global elasmobranch diversity and parasitology. *Bulletin of the American Museum of Natural History* **367**, 1–262. doi:10.1206/754.1
- NCBI Resource Coordinators** (2024) Database resources of the National Center for Biotechnology Information. Available at <https://www.ncbi.nlm.nih.gov/nucleotide/> (accessed 11 March 2024).
- Northeast Fisheries Science Center** (2006) *43rd Northeast Regional Stock Assessment Workshop (43rd SAW) assessment summary report*. NOAA Fisheries.
- Pacoureau N, Rigby CL, Kyne PM, Sherley RB, Winker H, Carlson JK, Fordham SV, Barreto R, Fernando D, Francis MP, Jabado RW, Herman KB, Liu K-M, Marshall AD, Pollom RA, Romanov EV, Simpfendorfer CA, Yin JS, Kindsvater HK and Dulvy NK** (2021) Half a century of global decline in oceanic sharks and rays. *Nature* **589**, 567–571. doi:10.1038/s41586-020-03173-9
- Parker S and Francis M** (2012) *Productivity of Two Species of Deepwater Sharks, Deania Calcea and Centrophorus Squamosus in New Zealand*. Ministry for Primary Industries: Wellington, New Zealand.
- Quiroz J-C, Wiff R, Gatica C and Leal E** (2008) Composición de especies, tasas de captura y estructura de tamaño de peces capturados en la pesquería espinelera artesanal de rayas en la zona sur-austral de Chile. *Latin American Journal of Aquatic Research* **36**, 15–24. doi:10.3856/vol36-issue1-fulltext-2
- Sagarese S, Frisk M, Cerrato R, Sosebee K, Musick J and Rago PJ** (2013) Application of generalized additive models to examine ontogenetic and seasonal distributions of Spiny Dogfish (*Squalus acanthias*) in the Northeast (US) shelf large marine ecosystem. *Canadian Journal of Fisheries and Aquatic Sciences* **71**, 847–877. doi:10.1139/cjfas-2013-0342
- Sánchez N, González HE and Iriarte JL** (2011) Trophic interactions of pelagic crustaceans in Comau Fjord (Chile): Their role in the food web structure. *Journal of Plankton Research* **33**, 1212–1229. doi:10.1093/plankt/fbr022
- Sims D** (2006) Differences in habitat selection and reproductive strategies of male and female sharks. *Sexual Segregation in Vertebrates: Ecology of the Two Sexes*. Cambridge, UK, pp. 127–147.
- Soto D, León-Muñoz J, Dresdner J, Luengo C, Tapia FJ and Garreaud R** (2019) Salmon farming vulnerability to climate change in southern Chile: Understanding the biophysical, socioeconomic and governance links. *Reviews in Aquaculture* **11**, 354–374. doi:10.1111/raq.12336
- Soto M-V, Arriagada-González J, Molina-Benavides M, Cabello M, Contreras-Alonso M, Ibarra I, Guevara G, Sepúlveda SA and Maerker M** (2023) Geodynamic ‘Hotspots’ in a periglacial landscape: Natural hazards and impacts on productive activities in Chilean Fjordlands, Northern Patagonia. *Geosciences* **13**.
- Spalding M, Fox H, Allen G, Davidson N, Ferdaña Z, Finlayson M, Halpern B, Jorge M, Lombana A, Lourie S, Martin K, McManus E, Molnar J, Recchia C and Robertson J** (2007) Marine ecoregions of the world: A bioregionalization of coastal and shelf areas. *BioScience* **57**, 573–583.
- Springer S** 1979 (Apr.) A revision of the catsharks, family Scyliorhinidae. NOAA (National Oceanic and Atmospheric Administration) Technical Report NMFS (National Marine Fisheries Service) Circular No. 422: i–v + 1–152.
- Stecher G, Tamura K and Kumar S** (2020) Molecular Evolutionary Genetics Analysis (MEGA) for macOS. *Molecular Biology and Evolution* **37**, 1237–1239.
- Stenberg C** (2005) Life history of the Piked Dogfish (*Squalus acanthias* L.) in Swedish waters. *Journal of Northwest Atlantic Fishery Science* **37**, 155–164.
- Straube N, White WT, Ho H-C, Rochel E, Corrigan S, Li C and Naylor GJP** (2013) A DNA sequence-based identification checklist for Taiwanese chondrichthyans. *Zootaxa* **3752**, 256–278.
- Sulikowski J, Prohaska B, Carlson A, Cicia A, Brown C and Morgan A** (2013) Observations of neonate spiny dogfish, *Squalus acanthias*, in Southern New England: A first account of a potential pupping ground in the Northwestern Atlantic. *Fisheries Research* **137**, 59–62.
- Templeman W** (1944) The life history of the spiny dogfish (*Squalus acanthias*) and the vitamin A. values of dogfish liver oil. *Research Bulletin No. 15 (Fisheries)* Department of Natural Resources, Newfoundland. St. John's, Newfoundland and Labrador: Memorial University of Newfoundland. Centre for Newfoundland Studies, p. 108.
- Thompson JD, Higgins DG and Gibson TJ** (1994) CLUSTAL W: Improving the sensitivity of progressive multiple sequence alignment through sequence

- weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Research* **22**, 4673–4680.
- Valenzuela A, Bustamante C and Lamilla J** (2008) Morphological characteristics of five bycatch sharks caught by southern Chilean demersal longline fisheries. *Scientia Marina* **72**, 231–237.
- Vargas-Caro C, Bustamante C, Bennett MB and Ovenden JR** (2017) Towards sustainable fishery management for skates in South America: The genetic population structure of *Zearaja chilensis* and *Dipturus trachyderma* (Chondrichthyes, Rajiformes) in the south-east Pacific Ocean. *PloS One* **12**, 1–12. doi:10.1371/journal.pone.0171111
- Villalobos VI, Valdivia N, Försterra G, Ballyram S, Espinoza JP, Wadham JL, Burgos-Andrade K and Häussermann V** (2021) Depth-dependent diversity patterns of rocky subtidal macrobenthic communities along a temperate Fjord in Northern Chilean Patagonia. *Frontiers in Marine Science* **8**, 1–18. doi:10.3389/fmars.2021.635855
- Williams G, Andrews K, Farrer D and Levin P** (2010) Catch rates and biological characteristics of Bluntnose Sixgill Sharks in Puget Sound. *Transactions of the American Fisheries Society* **139**, 108–116. doi:10.1111/j.1365-3113.2009.00431.x